

姓名: Sol

葉均承 應數一線性代數

學號: \_\_\_\_\_

**Quiz 1**

考試日期: 2020/09/24

1. 請框出答案. 2. 不可使用手機、計算器, 禁止作弊! 3. 背面還有題目

1. (50%) Determine whether the vector  $\vec{b}$  is in the span of the vectors  $\vec{v}_i$ . If so, write  $\vec{b}$  into the linear combination form.

$$\vec{b} = \begin{bmatrix} 3 \\ 5 \\ 3 \end{bmatrix}, \vec{v}_1 = \begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ 4 \\ -2 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} -3 \\ -1 \\ 5 \end{bmatrix}$$

$$\left[ \begin{array}{ccc|c} 0 & 1 & -3 & 3 \\ 2 & 4 & -1 & 5 \\ 4 & -2 & 5 & 3 \end{array} \right] \rightsquigarrow \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{array} \right]$$

$\therefore$  Yes!  $\vec{b} \in \text{sp}(\vec{v}_1, \vec{v}_2, \vec{v}_3)$

$$\vec{b} = 2\vec{v}_1 + 0\vec{v}_2 - \vec{v}_3$$

# Quiz 1

應數一線性代數

2. (50%) Prove that the given relation holds for all vectors, matrices and scalars for which the expression are defined.

$$(AB)^T = B^T A^T$$

$\triangle A = [a_{ij}]$ ,  $B = [b_{ij}]$ , let  $AB = C = [c_{ij}]$

Note  $c_{ij} = \sum_k a_{ik} b_{kj}$  :-( $AB)^T = C^T = [c'_{ij}] \therefore c'_{ij} = c_{ji} = \sum_k a_{jk} b_{ki}$

$\triangle A^T = [a'_{ij}]$ ,  $B^T = [b'_{ij}]$ , let  $B^T A^T = D = [d_{ij}]$

$$a'_{ij} = a_{ji} \quad b'_{ij} = b_{ji}$$

$$d_{ij} = \sum_k b'_{ik} a'_{kj} = \sum_k b_{ki} a_{jk} = \sum_k a_{jk} b_{ki} = c'_{ij}$$

$$\therefore C = D$$