

應數一線性代數 2022 春, 期中考 解答

學號: _____, 姓名: _____

本次考試共有 10 頁 (包含封面), 有 10 題。如有缺頁或漏題, 請立刻告知監考人員。

考試須知:

- 請在第一及最後一頁填上姓名學號, 並在每一頁的最上方屬名, 避免釘書針斷裂後考卷遺失。
- 不可翻閱課本或筆記。
- 計算題請寫出計算過程, 閱卷人員會視情況給予部份分數。
沒有計算過程, 就算回答正確答案也不會得到滿分。
答卷請清楚乾淨, 儘可能標記或是框出最終答案。

高師大校訓: 誠敬宏遠

誠, 一生動念都是誠實端正的。 敬, 就是對知識的認真尊重。
宏, 開拓視界, 恢宏心胸。 遠, 任重致遠, 不畏艱難。

請尊重自己也尊重其他同學, 考試時請勿東張西望交頭接耳。

1. (10 points) (a) Solve the system $\begin{cases} x_1' = 3x_1 - 5x_2 \\ x_2' = 2x_2 \end{cases}$
(b) Find the solution that satisfies the initial condition $x_1(0) = 2, x_2(0) = 5$.

Answer: (a) $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 5k_1e^{2t} + k_2e^{3t} \\ k_1e^{2t} \end{bmatrix}$, (b) $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 25e^{2t} - 23k_2e^{3t} \\ 5e^{2t} \end{bmatrix}$

Follow 課本 5-3 example 3

Follow 109-2 midterm problem 1.

2. (10 points) Let

$$A = \begin{bmatrix} 9 & -3 & 3 \\ -2 & 10 & 2 \\ 1 & 1 & 11 \end{bmatrix}$$

Find (if exists) an invertible matrix C and a diagonal matrix D such that $D = C^{-1}AC$. Also, find the eigenvalues of A^{100} .

(1) Is A diagonalizable? YES!. If A diagonalizable, $C = \begin{bmatrix} 0 & 1 & 3 \\ 1 & 0 & 2 \\ 1 & 1 & -1 \end{bmatrix}$, $D = \begin{bmatrix} 12 & 0 & 0 \\ 0 & 12 & 0 \\ 0 & 0 & 6 \end{bmatrix}$.

(2) The eigenvalue of A are 6, 12. The eigenvalue of A^{100} are $6^{100}, 12^{100}$.

Follow 課本 5-2 example 3, 4 and Theorem 5.1

Follow 108-2 midterm problem 1.

3. (10 points) Find the formula for the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that reflects in the line $3x + 2y = 0$.

Answer: $T([x, y]) = \underline{\frac{-1}{13}[5x + 2y, 12x - 5y]}$

Follow 課本 5-2 example 2

Follow 109-2 midterm problem 2.

4. (10 points) Find all the possible a, b, c, d, x, y so that the matrix A is orthogonal.

$$A = \begin{bmatrix} a & y & 0 \\ 2x & 3y & c \\ x & b & d \end{bmatrix}$$

Let

$$\vec{v}_1 = \begin{bmatrix} a \\ 2x \\ x \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} y \\ 3y \\ b \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 0 \\ c \\ d \end{bmatrix}$$

Since the matrix A is orthogonal, we have $0 = \vec{v}_1 \cdot \vec{v}_2 = \vec{v}_1 \cdot \vec{v}_3 = \vec{v}_2 \cdot \vec{v}_3$ and $1 = |\vec{v}_1|, |\vec{v}_2|, |\vec{v}_3|$ is zero vector.

In the case of $x = 0$. Since $1 = |\vec{v}_1|$, we have $a = 1$. Using $0 = \vec{v}_1 \cdot \vec{v}_2 = ay = y$, we have $y = 0$ and $b = 1$ ($1 = |\vec{v}_2|$). Thus $c = 1, d = 0$. Therefore, we have one solution $[a, b, c, d, x, y] = [1, 1, 1, 0, 0, 0]$

In the case of $x \neq 0$. By $0 = \vec{v}_1 \cdot \vec{v}_3$, we have $2cx + dx = 0$. Thus $c : d = 1 : -2$. Since $1 = |\vec{v}_3|$,

$$\vec{v}_3 = \begin{bmatrix} 0 \\ c \\ d \end{bmatrix} = \frac{\pm 1}{\sqrt{5}} \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}.$$

Since $0 = \vec{v}_2 \cdot \vec{v}_3$ and $c : d = 1 : -2$, $3y : b = 2 : 1$. Thus $b = \frac{3y}{2}$ and $\vec{v}_2 = \begin{bmatrix} y \\ 3y \\ 3y/2 \end{bmatrix} = \frac{\pm 1}{7} \begin{bmatrix} 2 \\ 6 \\ 3 \end{bmatrix}$.

Since $0 = \vec{v}_1 \cdot \vec{v}_2 = 2a + 12x + 3x$, we have $a = \frac{15x}{2}$. Thus $\vec{v}_1 = \begin{bmatrix} \frac{15x}{2} \\ 2x \\ x \end{bmatrix} = \frac{\pm 1}{\sqrt{245}} \begin{bmatrix} 15 \\ 4 \\ 2 \end{bmatrix}$.

ANSWER: Therefore, we have the solution:

$$\vec{v}_1 = \begin{bmatrix} a \\ 2x \\ x \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} y \\ 3y \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 0 \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

or

$$\vec{v}_1 = \begin{bmatrix} a \\ 2x \\ x \end{bmatrix} = \frac{\pm 1}{\sqrt{245}} \begin{bmatrix} 15 \\ 4 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} y \\ 3y \\ b \end{bmatrix} = \frac{\pm 1}{7} \begin{bmatrix} 2 \\ 6 \\ 3 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 0 \\ c \\ d \end{bmatrix} = \frac{\pm 1}{\sqrt{5}} \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}$$

5. (10 points) Find the projection matrix P for the plane $W : 2x - y + 2z = 0$ and then find the projection of $\vec{b} = [3, 2, 1]$ on the plane.

Answer: $\vec{b}_W = \underline{\frac{1}{3}[5, 8, -1]}$, $P = \underline{\frac{1}{9} \begin{bmatrix} 5 & 2 & -4 \\ 2 & 8 & 2 \\ -4 & 2 & 5 \end{bmatrix}}$.

Follow 課本 6-4 example 3

Follow 109-2 midterm problem 4.

6. (10 points) Let

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -1 \\ -1 & 1 & 1 \end{bmatrix}$$

Factor A in the form $A = QR$, where Q is an orthogonal matrix and R is an upper-triangular invertible matrix.

$$\text{Answer: } Q = \frac{1}{\sqrt{6}} \begin{bmatrix} \sqrt{3} & \sqrt{2} & 1 \\ 0 & \sqrt{2} & -2 \\ -\sqrt{3} & \sqrt{2} & 1 \end{bmatrix}, R = \begin{bmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{3} & \frac{-1}{\sqrt{3}} \\ 0 & 0 & \frac{\sqrt{8}}{\sqrt{3}} \end{bmatrix}.$$

Follow 課本 6-2 example 5.

Follow 109-2 midterm problem 6.

7. (10 points) Find the least squares straight line fit to the five points $(-4, -2)$, $(-2, 0)$, $(0, 1)$, $(2, 4)$, $(4, 5)$ and use it to approximate the fifth points $(1, a)$.

Answer: the line equation = $0.9x + 1.6$, $a = 2.5$.

Follow 課本 6-5 example 1

8. (10 points) Prove that, for every square matrix A all of whose eigenvalues are real, the product of its eigenvalues is $\det(A)$

Section 5-2, problem 17

If the characteristic polynomial of A is $p(\lambda) = |A - \lambda I|$, then $p(0) = |A| = \det(A)$.

Also,

$$p(\lambda) = (-1)^n (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n)$$

, so

$$p(0) = (-1)^{2n} \lambda_1 \lambda_2 \cdots \lambda_n = \lambda_1 \lambda_2 \cdots \lambda_n = \det(A).$$

9. (10 points) Show that the real eigenvalue of an orthogonal matrix must be equal to 1 or -1.

Hint: Think in terms of linear transformations.

Section 6-3, problem 27

