

1. 請框出答案. 2. 不可使用手機、計算器，禁止作弊!

1. Let the sequence  $a_0, a_1, a_2, \dots$  be given by  $a_0 = 2, a_1 = -1$  and  $a_k = \frac{1}{2}a_{k-1} + \frac{1}{2}a_{k-2}$  for  $k \geq 2$ .  
Find the explicit formula for  $a_k$ .

Answer:  $a_k = 0.5^{k-1}$

**Solution :**

$$\begin{bmatrix} a_{k+1} \\ a_k \end{bmatrix} = \begin{bmatrix} 0.5 & 0.5 \\ 1 & 0 \end{bmatrix}^k \begin{bmatrix} a_1 \\ a_0 \end{bmatrix}$$

Let

$$A = \begin{bmatrix} 0.5 & 0.5 \\ 1 & 0 \end{bmatrix}, \det(A - \lambda I) = (0.5 - \lambda)(-\lambda) - 0.5 = (1 - \lambda)(-0.5 - \lambda)$$

Therefore, the eigenvalues are  $\lambda_1 = 1, \lambda_2 = -0.5$ .

$\lambda_1 = 1$ ,

$$rref(A - \lambda_1 I) = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}, \vec{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$\lambda_2 = -0.5$ ,

$$rref(A - \lambda_2 I) = \begin{bmatrix} 1 & 0.5 \\ 0 & 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

**Method 1:**

Let

$$C = \begin{bmatrix} 1 & 1 \\ 1 & -2 \end{bmatrix}, D = \begin{bmatrix} 1 & 0 \\ 0 & 0.5 \end{bmatrix},$$

$$\begin{bmatrix} a_{k+1} \\ a_k \end{bmatrix} = A^k \begin{bmatrix} a_1 \\ a_0 \end{bmatrix} = (CDC^{-1})^k \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \frac{1}{2^k} \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

**Method 2:** Let

$$\begin{bmatrix} a_1 \\ a_0 \end{bmatrix} = r_1 \vec{v}_1 + r_2 \vec{v}_2$$

Then

$$\begin{bmatrix} a_{k+1} \\ a_k \end{bmatrix} = A^k \begin{bmatrix} a_1 \\ a_0 \end{bmatrix} = A^k(r_1 \vec{v}_1 + r_2 \vec{v}_2) = r_1 \lambda_1^k \vec{v}_1 + r_2 \lambda_2^k \vec{v}_2$$

Notice that  $r_1 = 0, r_2 = 1$

$$\begin{bmatrix} a_{k+1} \\ a_k \end{bmatrix} = A^k \begin{bmatrix} a_1 \\ a_0 \end{bmatrix} = 0.5^k \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -0.5^k \\ 0.5^{k-1} \end{bmatrix}$$

2. Solve the given system.

$$\begin{cases} x'_1 = x_1 - 3x_2 + 3x_3, \\ x'_2 = -5x_2 + 6x_3, \\ x'_3 = -3x_2 + 4x_3 \end{cases}$$

Answer:  $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 2 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} k_1 e^t \\ k_2 e^t \\ k_3 e^{-2t} \end{bmatrix} = \begin{bmatrix} k_1 e^t + k_3 e^{-2t} \\ k_2 e^t + 2k_3 e^{-2t} \\ k_2 e^t + k_3 e^{-2t} \end{bmatrix},$

---

**Solution :**

$$A = \begin{bmatrix} 1 & -3 & 3 \\ 0 & -5 & 6 \\ 0 & -3 & 4 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 2 \\ 0 & 1 & 1 \end{bmatrix}, D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix},$$

Such that  $AC = CD$ , and  $\vec{x}' = A\vec{x}$

Solve  $\vec{y}' = D\vec{y}$ , i.e.  $\begin{cases} y'_1 = y_1 \\ y'_2 = y_2 \\ y'_3 = -2y_3 \end{cases} \Rightarrow \begin{cases} y_1 = k_1 e^t \\ y_2 = k_2 e^t \\ y_3 = k_3 e^{-2t} \end{cases}$

$$\vec{x} = C\vec{y} \Rightarrow \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} k_1 e^t + k_3 e^{-2t} \\ k_2 e^t + 2k_3 e^{-2t} \\ k_2 e^t + k_3 e^{-2t} \end{bmatrix}$$