

1. 請框出答案. 2. 不可使用手機、計算器，禁止作弊!

1. Let $W = \text{sp}([0, 1, -1, -1], [1, -1, 1, 0], [-1, 1, 0, 1])$ is a subspace of \mathbb{R}^4 . Using the Gram-Schmidt process to find an orthonormal basis for W and then transform this to an orthonormal basis for \mathbb{R}^4 . Given $\vec{b} = [-1, 3, 2, 0]$, please find the projection \vec{b}_W .

Answer: an orthonormal basis for W is $\{\frac{1}{\sqrt{3}}[0, 1, -1, -1], \frac{1}{\sqrt{15}}[3, -1, 1, -2], \frac{1}{\sqrt{15}}[1, 3, 2, 1]\}$

$$\vec{b}_W = \underline{[0, 3, 1, 1]}$$

an orthonormal basis for \mathbb{R}^4 is $\{\frac{1}{\sqrt{3}}[0, 1, -1, -1], \frac{1}{\sqrt{15}}[3, -1, 1, -2], \frac{1}{\sqrt{15}}[1, 3, 2, 1], \frac{1}{\sqrt{3}}[1, 0, -1, 1]\}$

Solution :

Let $\vec{a}_1 = [0, 1, -1, -1]$, $\vec{a}_2 = [1, -1, 1, 0]$, $\vec{a}_3 = [-1, 1, 0, 1]$,

$$\vec{v}_1 = [0, 1, -1, -1],$$

$$\vec{q}_1 = \frac{\vec{v}_1}{\|\vec{v}_1\|} = \frac{1}{\sqrt{3}}[0, 1, -1, -1],$$

$$\vec{v}_2 = \vec{a}_2 - \frac{\vec{v}_1 \cdot \vec{a}_2}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 = \frac{1}{3}[3, -1, 1, -2],$$

$$\vec{q}_2 = \frac{\vec{v}_2}{\|\vec{v}_2\|} = \frac{1}{\sqrt{15}}[3, -1, 1, -2]$$

$$\vec{v}_3 = \vec{a}_3 - \frac{\vec{v}_1 \cdot \vec{a}_3}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 - \frac{\vec{v}_2 \cdot \vec{a}_3}{\vec{v}_2 \cdot \vec{v}_2} \vec{v}_2 = \frac{1}{5}[1, 3, 2, 1],$$

$$\vec{q}_3 = \frac{\vec{v}_3}{\|\vec{v}_3\|} = \frac{1}{\sqrt{15}}[1, 3, 2, 1],$$

Let $\vec{a}_4 = [1, 0, 0, 0]$, $\vec{a}_5 = [0, 1, 0, 0]$, $\vec{a}_6 = [0, 0, 1, 0]$, $\vec{a}_7 = [0, 0, 0, 1]$

$$\vec{v}_4 = \vec{a}_4 - \frac{\vec{v}_1 \cdot \vec{a}_4}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 - \frac{\vec{v}_2 \cdot \vec{a}_4}{\vec{v}_2 \cdot \vec{v}_2} \vec{v}_2 - \frac{\vec{v}_3 \cdot \vec{a}_4}{\vec{v}_3 \cdot \vec{v}_3} \vec{v}_3 = \frac{1}{3}[1, 0, -1, 1],$$

$$\vec{q}_4 = \frac{\vec{v}_4}{\|\vec{v}_4\|} = \frac{1}{\sqrt{3}}[1, 0, -1, 1],$$

Method 1:

$$\begin{aligned}\vec{b}_W &= (\vec{b} \cdot \vec{q}_1) \vec{q}_1 + (\vec{b} \cdot \vec{q}_2) \vec{q}_2 + (\vec{b} \cdot \vec{q}_3) \vec{q}_3 \\ &= [0, 3, 1, 1]\end{aligned}$$

Method 2:

$$\vec{b}_W = \vec{b} - \vec{b}_{W^\perp} = \vec{b} - (\vec{q}_4 \cdot \vec{b}) \vec{q}_4 = [0, 3, 1, 1]$$